

## ALLEGRA-D 12HR ALLERGY & CONGESTION TABLETS

### Drug Facts

#### Active ingredients (in each tablet)

Active ingredients (in each tablet)	Purpose
Fexofenadine HCl 60 mg	Antihistamine
Pseudoephedrine HCl 120 mg	Nasal decongestant

#### Uses

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
  - runny nose
  - itching of the nose or throat
  - sneezing
  - itchy, watery eyes
- temporarily relieves congestion due to the common cold, hay fever or other upper respiratory allergies
- reduces swelling of nasal passages
- temporarily relieves sinus congestion and pressure
- temporarily restores freer breathing through the nose

#### Warnings

##### Do not use

- if you have ever had an allergic reaction to this product or any of its ingredients
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product
- if you have difficulty swallowing

##### Ask a doctor before use if you have

- heart disease
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- kidney disease. Your doctor should determine if you need a different dose.
- thyroid disease
- diabetes
- glaucoma

#### When using this product

- do not take more than directed**
- do not take at the same time as aluminum or magnesium antacids
- do not take with fruit juices (see Directions)
- the tablet coating may be seen in the stool (this is normal). Continue to take as directed (see Directions).

#### Stop use and ask a doctor if

- an allergic reaction to this product occurs. Seek medical help right away.
- symptoms do not improve within 7 days or are accompanied by a fever
- you get nervous, dizzy, or sleepless

If pregnant or breast-feeding, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

#### Directions

- do not divide, crush, chew or dissolve the tablet; swallow tablet whole

adults and children 12 years of age and over	take 1 tablet with a glass of water every 12 hours on an empty stomach; do not take more than 2 tablets in 24 hours
children under 12 years of age	do not use
adults 65 years of age and older	ask a doctor
consumers with kidney disease	ask a doctor

#### Other information

- safety sealed: do not use if carton is opened or if individual blister units are torn or opened
- store between 20° and 25°C (68° and 77°F)

#### Inactive ingredients

carnauba wax, colloidal silicon dioxide, croscarmellose sodium, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch, stearic acid

#### Questions or comments?

call toll-free 1-800-633-1610 or [www.allegra.com](http://www.allegra.com)

[ALLEGRA DFL.]

